




TECHNOLOGY

Use of Biomarkers for Assessing Differential Risk for Developing Heart Failure Phenotypes

OVERVIEW

Left ventricular (LV) hypertrophy (LVH), most commonly due to chronic hypertensive heart disease, is associated with increased risk of developing depressed LV systolic function, heart failure (HF) and death. Routine cardiac imaging to screen for LVH in hypertensive patients is not recommended because of the low predictive value of LVH for HF. Researchers at UMB have demonstrated that the presence of elevated cardiac-specific biomarkers in the presence of LVH identifies subjects with a high risk of developing HF, particularly HF with reduced ejection fraction (HFrEF). This finding is significant because it identifies patients that would benefit from specific medical and lifestyle interventions.

UMB researchers have shown that in individuals aged 65 or older with LVH and a significant, sustained increase in NT-proBNP or hs-cTnT levels have markedly higher rates of incident HF, in particular, HFrEF. Specifically, a significant, sustained increase in the N-terminal pro-b-type natriuretic peptide (NT-proBNP) and high-sensitivity cardiac troponin T (hs-CTnT) in patients exhibiting LVH, there was a 3-fold increase in the development of HFrEF. HFrEF occurs in approximately half of the patients that develop HF and is the only form of HF that is currently treatable. This discovery allows for the first time a diagnostic to identify patients with a high risk of developing HF. Predicting HF provides, for the first time, an opportunity to modify the treatment and lifestyle of high-risk patients to prevent disease.



APPLICATIONS

The prevalence of LVH in the general population ranges from 9.1% for women and 14.9% for men, or approximately 35 million people in the US. Heart failure is considered an emerging epidemic, with a US prevalence greater than 5.8 million people. Roughly half of all people who develop heart failure die within 5 years of diagnosis. Echocardiogram is used to establish LVH, but cannot reliably predict which patients will develop heart failure.

ADVANTAGES

Predicts the progression to HF following asymptomatic LVH

Specifically associated with HFpEF, which can guide therapeutic interventions

Biomarkers may be developed as therapeutic targets

STAGE OF DEVELOPMENT

This technology has been tested in a clinical samples from the general population and from adults without a prior history of heart failure or myocardial infarction. Additional clinical studies are ongoing.

R&D REQUIRED

The prevalence of LVH in the general population ranges from 9.1% for women and 14.9% for men, or approximately 35 million people in the US. HF is considered an emerging epidemic, with a US prevalence greater than 5.8 million people. There is currently no diagnostic that can be used to predict the risk of patients with LVH developing HF.

CONTACT INFO

Office of Technology Transfer
620 W Lexington St., 4th Floor
Baltimore, MD 21201
Email: ott@umaryland.edu
Phone: (410) 706-2380

Additional Information

INSTITUTION

University of Maryland, Baltimore

PATENT STATUS

US 10,509,044

LICENSE STATUS

Available for licensing

CATEGORIES

- Diagnostics
- Biomarker

INVESTIGATOR(S)

Christopher deFilippi
James de Lemos
Robert H. Christenson
Stephen L. Seliger

ATTACHMENTS

-  [Download CD-2014-098 Marketing 10_16_2019 FINAL.pdf](#)

EXTERNAL RESOURCES

- [High Sensitive Cardiac Troponin T as an Early Biochemical Signature for Clinical and Subclinical Heart Failure: The Multi-Ethnic](#)
- [CHS \(Cardiovascular Health Study\)](#)
- [Biomarkers of chronic cardiac injury and hemodynamic stress identify a malignant phenotype of LVH in the general population](#)

CD-2014-098