



## TECHNOLOGY

# Stable Fluorescent Mutant of *Perkinsus marinus* Expressing GFP

## OVERVIEW

*Perkinsus* are intracellular parasites that cause disease in shellfish, especially bivalve molluscs. Its infection has devastated oyster populations along the US coasts. In addition to the economic value of shellfisheries, filter-feeding bivalves such as oysters play a critical role in maintaining water quality, making *Perkinsus* a serious threat to the economy and the health and integrity of the coastal ecosystem. Numerous attempts to control Dermo disease, a result of *P. marinus* infection, have been unsuccessful to date, partly due to a lack of information about fundamental aspects of *P. marinus* biology. UMB researchers have developed a transfection system for Perkinsozoa and created a stable, fluorescent mutant of *P. marinus* that expresses GFP. This transgenic organism can be used to study the mechanisms underlying *P. marinus* infection in bivalves, allowing for easier visualization and quantification.

Using optimized *P. marinus* culture methods, UMB researchers constructed a vector based on a highly expressed *P. marinus* gene tagged with GFP (pPmMOE-GFP). Under optimized transfection conditions, exogenous pPmMOE-GFP DNA was introduced into *P. marinus* trophozoites by electroporation with a robust efficiency of transfection of ~38%. Fluorescence can be detected starting 14 hours after electroporation and stably expressed for longer than 8 months. The visualization allows researchers to study its location, growth rate, and proliferation.

## APPLICATIONS

Bivalves such as oysters are important both ecologically and economically. As the causing agent of Dermo disease in oysters, *P. marinus* has caused mass mortality of Eastern oysters commonly cultured and wild-harvested as food species. Found along the eastern coast of the USA, from Maine to Florida, the presence of this parasite has severely reduced the abundance and productivity of oysters, particularly that of *Crassostrea virginica*. In the Mid-Atlantic region, *P. marinus* is responsible for a decrease of at least two-thirds of the surplus production available to fisheries. GFP-labeled *P. marinus* can help better understand the biology behind its infection and assist in identifying successful intervention strategies to curb it, both crucial to protecting the ecosystem and aquaculture industries.

## ADVANTAGES

GFP-tagged *P. marinus* allows easier visualization and quantification of infection in bivalves

GFP-expression is induced early and stably, lasting longer than 8 months

## STAGE OF DEVELOPMENT

pPmMOE-GFP-tagged *P. marinus* has been tested *in vivo*, where PmMOE-GFP is expressed with a strong concentration in the inner lining of the plasma membrane.

(As of 5/2/17)

## LICENSING POTENTIAL

Available for non-exclusive licensing

## **CONTACT INFO**

Office of Technology Transfer  
620 W Lexington St., 4th Floor  
Baltimore, MD 21201  
Email: [ott@umaryland.edu](mailto:ott@umaryland.edu)  
Phone: (410) 706-2380

## **Additional Information**

### **INSTITUTION**

University of Maryland, Baltimore

### **LICENSE STATUS**

A non-exclusive license is available for this technology

### **CATEGORIES**

- Research Tools, Antibodies, & Reagents

### **INVESTIGATOR(S)**

Gerardo R. Vasta  
Jose-Antonio Fernandez-Robledo

### **ATTACHMENTS**

-  [Download GV-2014-015 Marketing Sheet 5-2-17 FINAL.pdf](#)

### **EXTERNAL RESOURCES**

- [Transfection of the protozoan parasite Perkinsus marinus.](#)

GV-2014-015